

|                   |   |
|-------------------|---|
| Postprint Version | 1.0   |
| Journal website   | <a href="http://jech.bmj.com/cgi/content/abstract/jech.2008.079038v1">http://jech.bmj.com/cgi/content/abstract/jech.2008.079038v1</a> |
| Pubmed link       | <a href="http://www.ncbi.nlm.nih.gov/pubmed/19833605">http://www.ncbi.nlm.nih.gov/pubmed/19833605</a>                                 |
| DOI               | 10.1136/jech.2008.079038  |

This is a NIVEL certified Post Print, more info at <http://www.nivel.eu>

## Morbidity is related to a green living environment

JOLANDA MAAS M.Sc. <sup>(1)</sup>, ROBERT A. VERHEIJ PHD <sup>(1)</sup>, SJERP DE VRIES PHD <sup>(3)</sup>, PETER SPREEUWENBERG <sup>(1)</sup>, FRANCOIS G. SCHELLEVIS MD, PHD <sup>(1)(4)</sup>, PETER P. GROENEWEGEN PHD <sup>(1)(2)</sup>

<sup>1)</sup> NIVEL (Netherlands Institute for Health Services Research), Utrecht, the Netherlands

<sup>2)</sup> Utrecht University, Department of Human Geography, Department of Sociology, P.O. Box 80115, NL-3508 TC Utrecht, The Netherlands

<sup>3)</sup> Alterra, Green World Research, P.O. Box 47 NL-6700 AA Wageningen, The Netherlands

<sup>4)</sup> Department of General Practice/ EMGO Institute VU University Medical Centre, Amsterdam, The Netherlands

Correspondence to: J. Maas MSc.,  
NIVEL, Netherlands Institute for Health Services Research  
P.O Box 1568 3500 BN Utrecht the Netherlands  
Tel: +31-30-2729846  
Fax: +31-30-2729729  
e-mail address: [j.maas@nivel.nl](mailto:j.maas@nivel.nl)

### ABSTRACT

**Background** Due to increasing urbanisation people face the prospect of living in environments with few green spaces. There is increasing evidence for a positive relation between green space in people's living environment and self-reported indicators of physical and mental health. This study investigates whether physician assessed morbidity is also related to green space in people's living environment.

**Methods** Morbidity data were derived from electronic medical records of 195 general practitioners in 96 Dutch practices, serving a population of 345,143 people. Morbidity was classified by the general practitioners according to the International Classification of Primary Care (ICPC). The percentage of green space within a one kilometre and three kilometre radius around the postal code coordinates was derived from an existing database and was calculated for each household. Multilevel logistic regression analyses were performed controlling for demographic and socio-economic characteristics.

**Results** The annual prevalence rate of 15 of the 24 disease clusters was lower in living environments with more green space in a 1 km radius. The relation was strongest for anxiety disorder and depression. The relation was stronger for children and people with a lower socioeconomic status. Furthermore, the relation was strongest in slightly urban areas and not apparent in very strongly urban areas.

**Conclusion** This study indicates that the previously established relation between green space and a number of self-reported general indicators of physical and mental health can also be found for clusters of specific physician assessed morbidity. The study stresses the importance of green space close to home for children and lower socio-economic groups.

## INTRODUCTION

Due to increasing urbanisation, combined with a planning policy of spatial densification, more people face the prospect of living in residential environments with little green space. At the same time, increasing evidence shows that green space has beneficial effects on people's health. Evidence has been found for a positive relation between green space and self perceived health, [1-4], longevity, [5], number of symptoms and the risk of psychiatric morbidity.[1] Access to a garden and shorter distances to green areas from the dwelling were associated with less stress and a lower likelihood of obesity. [6] Experimental studies showed that there is a positive relation between green space and restoration from stress and mental fatigue. More specific, exposure to nature has been found to have a positive effect on mood, concentration, self-discipline, and physiological stress. [7-11] These studies indicate that there is a relation between green space and self reported general indicators of physical and mental health. Thus, people living in greener environments report a better physical and mental health.

The decrease of green space could therefore have health consequences. However, it remains unknown whether living in residential environments with little green space also has negative consequences for objective health. In this explorative study we will go one step further than other studies and investigate whether the prevalence of several physician assessed morbidity clusters is also related to the amount of green space in people's living environment. This is the first study to investigate the relation between green space and prevalence of physician assessed morbidity. This study has an explorative character and takes into account a broad number of diseases highly prevalent in society.

To gain more insight into the relation between green space and physician assessed morbidity we analysed this relation separately for different age groups and different socio-economic groups. We hypothesise that the relation is stronger for elderly people and children (as compared to adults) because, as a result of their lower mobility, they spend more time in the vicinity of their home, resulting in higher exposure to green space in their living environment.

The same applies to people with a lower socio-economic status, whose activities and social contacts are situated close to their homes. [12,13] Therefore we also hypothesise that people with a lower socio-economic status are more exposed to the green space in their living environment. Finally, the relation was analysed for different levels of urbanicity to investigate whether the relation varies between urban and rural areas.

## METHODS

For this study data from two different datasets were combined. Morbidity data were collected within the framework of the second Dutch National Survey in General Practice (DNSGP-2), which included a nationwide, representative sample of 104 general practices with 195 GPs and a practice population of approximately 400 000 enlisted people, who were representative for the Dutch population in terms of age, gender and type of health insurance.[14] For this study data from 96 practices that recorded morbidity for a full period of 12 months or more were used. This selection had no significant effect on the representativeness of the data, because after the selection the sample was still representative for the Dutch population. [14] Only people who had been registered with their current GP for longer than 12 months prior to the study ( $n = 345\ 143$ ) were included, because we assumed that people will have to live for at least 12 months in the same living environment before any effect of it would be noticeable.

Environmental data were derived from the National Land Cover Classification database (LGN4) in 2001, which contains the dominant type of land use of each 25 x 25 metre grid cell of the Netherlands.[15] The two datasets were matched on the basis of the x and y coordinates of the respondent's six character postal code (on average about 15 to 20 households have the same six character postal code). The dataset included 50187 postal codes and on average 6.9 respondents resided in each postal code area.

### Morbidity data

Morbidity data were derived from routine primary care electronic medical records. In the Netherlands morbidity presented in general practice is a good indicator of morbidity in the population [16]. Basically all non-institutionalized people are registered with a GP.

Furthermore, GPs have a gate keeping role for secondary care and are usually the first point of contact with the health care system. The data have been validated for obtaining prevalence estimates [17] During a of 12 months period, data on all GP consultations with patients were extracted from the electronic medical records. These data included contact diagnoses and indications (diagnoses) for medication and referral to

secondary care. Prevalence rates are based on contacts that were classified by the GP according to the International Classification of Primary Care (ICPC) and subsequently clustered into episodes of disease. [18] The most prevalent episodes were combined into 24 disease clusters. These disease clusters have been used in several other studies [19, 20] and include the most prevalent diseases in general practice (prevalence > 10 per 1000) (see table 1). The 24 disease clusters have been distributed over disease categories, namely cardiovascular diseases, musculoskeletal diseases, mental diseases, respiratory diseases, neurological diseases, digestive diseases and miscellaneous.

Not all disease clusters were relevant for all age groups, therefore the epidemiological denominator varied (table 1). A prevalence rate for each cluster was calculated by dividing the number of patients with at least one disease episode in 2001 belonging to the cluster by the population at risk. The population at risk was based on age groups in which the diseases occurred. Some disease clusters, like for instance high blood pressure, were only present in the older age-groups. Therefore, for high blood pressure the prevalence rate was calculated by dividing the number of patients with at least one high blood pressure episode in 2001 by the older age-groups and not for the population as a whole.

#### [TABLE 1]

##### **Characteristics of the respondents' living environment**

The LGN4 database discriminates 39 land use classes including crop types, forest types, water, various urban classes and semi-natural classes and has been proven to be valid and accurate. [15, 21] The total percentage of green space in the respondents' living environment was measured within a 1-km radius and within a 3-km radius around the postcode centroid of a respondent's home, to see whether there is a stronger relation for green space close by than green space further away. Only green spaces that dominate the land use in the 25 by 25 meter grid cell (more than 50% of the grid cell is green) have been classified as green space in the dataset. Small-scale green spaces, such as street trees and roadside vegetation were only included as green space if they were dominant in the grid cell.

##### **Urbanicity**

Another environmental characteristic is urbanicity. This variable consists of five categories ranging from very strongly urban (1) to non-urban (5); it was measured at municipal level and was derived from Statistics Netherlands. The indicator is based on the number of households per square km and is commonly used in the Netherlands. [22]

##### **Demographic and socioeconomic characteristics**

Part of the relation between green space and health may be the result of direct or indirect selection. Direct selection would take place when people's health is related to their chances of living in a green environment. Indirect selection takes place when people with certain characteristics related to wellbeing (such as income) tend to live in a green environment [23].

As migration flows are related to such socio-demographic characteristics as age, income and education [24], we decided to rule out indirect selection effects as far as possible by controlling statistically for demographic and socio-economic characteristics.

The demographic characteristics taken into account were gender (female = 1) and age (which was taken into account as polynomial till the third order because there was no linear relation between the disease clusters and age) and were derived from the patient lists of the participating practices. To find out whether the relation between green space and morbidity differed between age groups, age was divided into six categories (viz. children (aged <12 year), adolescent (aged 13-17 year), youth (aged 18-25 year), young adults (aged 26-45 year), older adults (aged 46-65 year) and elderly (aged 65+)).

Socio-economic characteristics were collected by a registration form that was sent by mail to all people listed in the participating practices in the DNSGP-2 (n= 380 000, response 76,5%) [14] and included education, work status, and health care insurance type. For a number of people these socio-economic characteristics were unknown. To reduce the number of missings we included a category unknown in the analyses. Education was measured as the highest level of completed education (unknown, no education completed, primary education, secondary education, higher education). Work situation was categorised as: work situation unknown, paid job, attending school/ studying, housewife/ houseman, retired, disability pension, unemployed. Socio-economic status was additionally implicitly measured by type of health care insurance (unknown, public or private). The type of health care insurance can be regarded as an indicator of

socio-economic status in the Dutch context in 2001, as only people with a higher income had private health insurance, whereas people with a lower income had obligatory public health insurance.

When testing the effect of green space for different SES groups, SES was operationalised as the level of education divided into three categories, viz. higher education (university or higher vocational education), secondary education and primary or no education. Characteristics of the study population are displayed in table 2.

#### [TABLE 2]

##### **Statistical analysis**

The relation between percentage of green space in people's living environment and morbidity was assessed using multilevel logistic regression analyses, controlling for demographic and socio-economic characteristics and urbanicity. We included two levels: individuals and practices, because of the hierarchical structure of the data within DNSGP-2. The multilevel logistic regression analyses were performed with MLwiN. The independent variables, including the percentage of green space, were centred around their average. The results thus represent morbidity of the average population living in an area with an average amount of green space. We used interaction effects between respective age groups, SES groups and urbanicity and the green space indicator to investigate the relation for different age groups, SES groups and in different levels of urbanicity. Because of the large dataset we adopted a strict type 1 error criterion of  $\alpha = .01$ .

#### **RESULTS**

On average there is 42.4% of green space in a 1 km radius and 60.8% of green space in a 3km radius around people's home. Table 3 presents the odds ratios for the annual prevalence rate of the 24 disease clusters for people who have 10 percent more green space than average. In general, a significant relation between the percentage of green space and the annual prevalence rate was only present for green space in a 1 km radius. Only for anxiety disorders, infectious diseases of the digestive system and Medically Unexplained Physical Symptoms (MUUPS) the annual prevalence rate was lower in environments with more green space in a 3 km radius.

For fifteen of the twenty-four disease clusters the annual prevalence rate was lower in living environments with a higher percentage of green space in a 1km radius. This relation is apparent for diseases in all seven disease categories. It is strongest for anxiety disorders and depression. For none of the disease clusters the relationship is negative.

#### [TABLE 3]

##### **Strength of the relation**

An indication of the strength of the relation is given in table 4 which shows the annual prevalence per 1000 for people with average characteristics on the control variables with respectively 10% and 90% green space in a 1 km radius around their home. For anxiety disorders, the annual prevalence for people with average characteristics with 10% green space in a 1 km radius was 26 per 1000 people and for those with 90% green space in a 1 km radius 18 per 1000 people. For depression these figures are respectively 32 and 24 per 1000.

Generally the found relation between green space and physician assessed morbidity is comparable with the relation between age and morbidity. An increase of one percentage of green space on physician assessed morbidity equals the effect of one year lower age.

#### [TABLE 4]

##### **Relation in different age groups**

Further analysis showed that the relation was strongest for children younger than 12 and people between 46 and 65 (not in table). For children the relation was not only apparent for the percentage of green space in a 1 km radius, but also for the percentage of green space in a 14 3 km radius. For a few disease clusters the relation for children was especially strong, for example for vertigo (1 km: OR = 0.81 (95% C. I = 0.74 - 0.90) / 3km: OR 0.85 (95% C. I = 0.77 -0.94)) and severe intestinal complaints (1km: 0.85 (95% C. I = 0.80

- 0.90)/ 3km: 0.89 (95% C. I = 0.84 - 0.94)). The strongest relation for children was found for depression (1km: OR = 0.79 (95% C. I = 0.72 - 0.88) / 3 km: OR = 0.84 (95% C. I = 0.78 - 0.91)).

The relations for the other age groups were similar to the overall relations shown in table 3.

#### **Relation for different socio-economic groups**

Especially the lower educated groups had a lower annual prevalence rate when they had more green space in a 1 km radius around their home. For example, the odds for COPD were smaller for the lower educated (1 km: OR = .97 (95% C.I.= .95 - .99)) than for higher educated (OR = .98 (95% C.I.= .96 – 1.00)).

#### **Relation for different levels of urbanicity**

Concerning the level of urbanicity our analyses show that urbanicity influences the relation between green space and the annual prevalence of disease clusters (not in table). There is often no relation between green space and the annual prevalence of disease clusters in the very strongly urban areas. At all other levels of urbanicity people with more green space in a 1 km radius around their home had a lower annual prevalence rate. The relations between green space and annual prevalence rates were strongest in slightly urban areas.

## **DISCUSSION**

### **Principal findings**

This explorative study shows that the previously established relation between green space and a number of self-reported general indicators of physical and mental health can also be found for specific, doctor-assessed disease categories. The annual prevalence rates for 15 of the 24 investigated disease clusters is lower in living environments with more green space in a 1 km radius. Green space close to home appeared to be more important than green space further away. This is in contrast with our previous studies [1, 2] which found the relation between self-perceived health and the amount of green space in a 1km and a 3km radius around people's home to be equally strong. It appears that for the prevalence of these more specific diseases green space close to home is more important. This study differs from other studies which mainly focussed on the relation between green space and self-perceived measures of physical and mental health [1,2,3,4,6]. This is the first study to assess the relation between green space and specific diseases which were derived from electronic medical records of GP's. This dataset helps better establish the relation between green space and health, because it used physician assessed morbidity as outcome, because there was no single source bias in the data, and because we used a large dataset which was representative for the Netherlands.

In line with our hypothesis the relation was strongest for people who were expected to spend more time in the vicinity of their homes, namely children and people with a lower socioeconomic status. However, contrary to our expectations the relation appeared to be stronger for people aged between 46 and 65 than for elderly. Concerning urbanicity, the relation appeared to be strongest in slightly urban areas. In very strongly urban areas there was no relation with the annual prevalence of disease clusters. This may be related to the fact that green spaces in highly urban areas are more often found to evoke feelings of insecurity [25], and thereby inhibiting their use. This study only gives some indications for the relation between green space and morbidity for different subgroups. Further research should focus specifically on one of the subgroups to investigate the relation for subgroups more thoroughly.

### *Underlying mechanisms*

The results of this study give some indications for the possible mechanism behind the relation between green space and health. Several mechanisms could be responsible for the relation between green space and health, of which the following are most commonly mentioned: recovery from stress and attention fatigue, encouragement of physical activity, facilitation of social contact and better air quality. [7, 26] What do the results tell us about the mechanism at work? The strong relation we particularly found for anxiety disorder and depression suggests that recovery from stress and attention fatigue might be the most likely mechanism behind the relation between green space and health, though facilitation of social contacts might also contribute. However, there is no reason to discard any of the other possible mechanisms. In living environments with more green space, the prevalence of most respiratory illnesses was lower, indicating that air quality could also be a possible mechanism behind the relation between green space and health. For diseases related to physical activity (diabetes, high blood pressure, musculoskeletal diseases) somewhat less strong relations were found. But as the associations were present, physical activity could also be a possible

mechanism. Further research will have to shed more light on the mechanisms behind the relation between green space and health.

This study shows that the role of green space in the living environment for health should not be underestimated. Most of the diseases which were found to be related to the percentage of green space in the living environment are highly prevalent in society and in many countries they are subject of large scale prevention programs. Furthermore, in many countries, diseases of the circulatory system, mental disorders and diseases of digestive system are among the most expensive diseases with respect to health care costs. [27] Our study contributes to the evidence that green space can help fight some major public health threats in western societies and should be allocated a more central position in spatial planning. Healthy planning should include a place for green space and policy makers should take the amount of green space in the living environment into account when endeavouring to improve the health situation of children and lower socio-economic groups.

### **Strength and limitations**

This is the first large epidemiological study investigating the relation between the amount of green space in the living environment of people and the prevalence of physician assessed morbidity. Morbidity data were derived from a different database than the data on green space; consequently, there is no single source or method bias. On the other hand, we don't have information on exposure time.

The morbidity data are accurate because they were extracted from routine electronic medical records of general practices, and the inter-observer reliability of grouping contacts into episodes was high. [16] The registration covered a 12-month period for each practice in order to eliminate seasonal influences. Considering the representativeness of the participating GPs and their patients – and the high validity of the data – the results of the present study can be assumed to validly represent morbidity in Dutch general practice.

Furthermore, because general practice in the Netherlands is usually the first point of contact with the health care system, and because the GP has a gatekeeping role for specialist care, and because there are no large geographic [28] or social differences in access to general practice, morbidity presented in general practice can be regarded as a very close approximation of morbidity present in the open population.

The data used for this study also have some shortcomings. First, our data on green space, although assessed on a small scale, does not take small green spaces in the living environment into account. A 25 by 25 meter grid cell was only regarded as green space when green space dominates in the grid cell. Gardens and small-scale green spaces, such as street trees and green verges which could also influence people's health, are not regarded as green space in our study. Consequently the effect of green space might be slightly underestimated.

Second, because of the cross-sectional design of the study, it is not possible to make strong inferences about the causality of the relations that were found. The observed effects of green space on health may partly be caused by selection. We tried to rule out this possibility by taking socio-economic and demographic characteristics into account, but the effects of selection cannot be ruled out completely. The results from the subgroup analyses by SES groups, however, make it rather unlikely that selection is the responsible mechanism. The relationship observed between green space and morbidity was stronger for the less well-educated group and this is exactly the subgroup that has fewer options in their choice of neighbourhood of residence. Our results may be influenced by selective migration based on people's health (direct selection). However, longitudinal studies on health related migration show that direct selection can not be held responsible for geographical differences that remain if socioeconomic and demographic factors are taken into account. [29, 30] Third, we tried to control as much as possible for individual socio-economic status. However, we did not have any information on the income of the respondents, which is a relevant indicator for socio-economic status. Furthermore, we did not control for other confounders at neighbourhood level, although different studies have shown that for example neighbourhood SES could also influence health [31, 32]. Because this was an explorative study we chose to keep the design somewhat easy. Further research should try to find out whether a relation can also be found when neighbourhood SES is controlled for.

### **ACKNOWLEDGEMENT**

This study was supported by a grant from the Netherlands Organisation for Scientific Research.

## COMPETING INTEREST

No conflicts of interest are reported by any of the authors.

## LICENSE FOR PUBLICATION STATEMENT

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in JECH and any other BMJPG products and sublicences such use and exploit all subsidiary rights, as set out in our licence (<http://jech.bmj.com/ifora/licence.pdf>).

### What is already known on this subject?

There is increasing evidence for a positive relation between green space in the living environment and a number of self-reported indicators of physical and mental health. Smallscale psychological research showed that exposure to green space has a positive effect on stress reduction and attention restoration. Several epidemiological studies have shown that green space is positively correlated with self-perceived health, number of symptoms experienced and mortality.

### What does this study add?

This study uses large scale representative medical record data to investigate whether the prevalence of a number of disease clusters is related to the amount of green space in people's living environment. The annual prevalence rates for 15 of the 24 investigated disease clusters is lower in living environments with more green space in a 1km radius around people's homes. The study stresses the importance of green space close to people's home. The relationship is particularly strong for children and lower socio-economic groups.

## REFERENCES

1. de Vries S, Verheij RA, Groenewegen PP, Spreeuwenberg P. Natural environments healthy environments? An exploratory analysis of the relationship between greenspace and health. *Environment and Planning A* 2003, 35: 1717-1731.
2. Maas J, Verheij RA, Groenewegen PP, de Vries S, Spreeuwenberg P. Green space, urbanity and health . how strong is the relation? *JECH* 2006, 60: 587-592.
3. Mitchell R, Popham F. Greenspace, urbanity and health: relationships in England. *JECH* 2007, 61: 681-683.
4. Sugiyama T, Leslie E, Giles-Corti B, Owen N. Associations of neighbourhood greenness with physical and mental health: do walking, social coherence and local social interaction explain the relationships?. *JECH* 2008, 62: 9-doi:10.1136/jech.2007.064287
5. Takano T, Nakamura K, Watanabe M. Urban residential environments and senior citizens' longevity in megacity areas. The importance of walkable green spaces. *JECH* 2002, 56: 913-918.
6. Nielsen TS, Hansen KB. Do green areas affect health? Results from a Danish survey on the use of green areas and health indicators. *Health & Place* 2007, 13: 839-850.
7. Health Council of the Netherlands, Dutch Advisory Council for research on Spatial Planning Nature and the Environment. *Nature and health. The influence of nature on social, psychological and physical well-being*. The Hague: Health Council of the Netherlands; RMNO; 2004.
8. van den Berg AE, Hartig T, Staats H. Preference for nature in urbanized societies: stress, restoration, and the pursuit of sustainability. *JSI* 2007.
9. Hartig T, Mang M, Evans GW. Restorative effects of natural environment experiences. *Environment and Behavior* 1991, 23: 3-27.
10. Kaplan R, Kaplan S. *The experience of nature. A psychological perspective*. Cambridge: Cambridge University Press.; 1989.
11. Ulrich R. View through a window may influence recovery from surgery. *Science* 1984, 224: 420-421.
12. Schwanen T, Dijst M, Dieleman FM. A microlevel analysis of residential context and travel time. *Environment and Planning A* 2002, 34: 1487-1507.
13. Harms L. *Op weg in de vrije tijd: context, kenmerken en dynamiek van vrijetijds mobiliteit*. Den Haag: Sociaal Cultureel Planbureau; 2006.
14. Westert GP, Schellevis FG, de Bakker D.H., Groenewegen PP, Bensing JM, van der Zee J. Monitoring health inequalities through general practice: the Second Dutch National Survey of General Practice. *European Journal of Public Health* 2005, 15: 59-65.

**TABLES**

Table 1: Annual prevalence rates of clusters of diseases presented in general practice (cases per 1000) (n = 345 143; unless stated otherwise)

| Cluster  | ICPC codes   | N (abs) | /1000 |
|--|--|---------|-------|
| <b><i>Cardiovascular</i></b>                   |  |         |       |
| High blood pressure (n = 273 925)              | K85 K86 K87  | 24778   | 90.5  |
| Cardiac disease                                | K71 K73 K74 K77 K78 K79 K80 K81 K82<br>K83 K84   | 9044    | 26.2  |
| Coronary heart disease (n = 240 825)           | K74 K75 K76  | 5804    | 24.1  |
| Stroke, brain hemorrhage (n = 240 825)         | K89 K90  | 2549    | 10.6  |
| <b><i>Musculoskeletal</i></b>                  |  |         |       |
| Neck- and back complaints                      | L01 L02 L03 L84 L86  | 32346   | 93.7  |
| Severe back complaints                         | L02 L03 L85 L86  | 25230   | 73.1  |
| Severe neck and shoulder complaints            | L01 L08 L83 L92  | 21236   | 61.5  |
| Severe elbow, wrist and hand complaints        | L10 L11 L12 L72 L74  | 7698    | 22.3  |
| Osteoarthritis (n = 240 825)                   | L89 L90 L91  | 4521    | 18.8  |
| Arthritis (n = 240 825)                        | L88 T92  | 3170    | 13.2  |
| <b><i>Mental</i></b>                           |  |         |       |
| Depression                                     | P03 P76  | 8859    | 25.7  |
| Anxiety disorder                               | P01 P74  | 8033    | 23.3  |
| <b><i>Respiratory</i></b>                      |  |         |       |
| Upper respiratory tract infection              | A77 R72 R74 R75 R76 R80  | 31457   | 91.1  |
| Bronchi(oli)tis / pneumonia                    | R78 R81  | 10806   | 31.3  |
| Asthma, COPD                                   | R91 R95 R96  | 12813   | 37.1  |
| <b><i>Neurological</i></b>                     |  |         |       |
| Migraine/severe headache                       | N01 N02 N03 N89 N90 N92  | 10629   | 30.8  |
| Vertigo  | N17  | 4023    | 11.7  |
| <b><i>Digestive</i></b>                        |  |         |       |
| Severe intestinal complaints                   | D81 D85 D86 D92 D93 D94  | 5264    | 15.3  |
| Infectious disease of the intestinal canal     | D70 D73  | 3816    | 11.1  |
| <b><i>Miscellaneous</i></b>                    |  |         |       |
| Medically Unexplained Physical Symptoms (MUPS) | A01 A04 D01 D08 D09 D12 D18 D21 D93<br>K01 K02 K04 L01 L02 L03 L08 L09 L14<br>L20 N01 N02 N17 P06 P20 R02 R21 T03<br>T07 T08     | 75774   | 219.5 |
| Chronical eczema                               | S86 S87 S88  | 22303   | 64.6  |
| Acute urinary tract infection                  | U70 U71 U72  | 13303   | 38.5  |
| Diabetes (n = 290 479)                         | T88 T90  | 9260    | 31.9  |
| Cancer   | A79 B72 B73 B74 D74 D75 D76 D77 F74<br>H75 K72 L71 N74 R84 R85 S77 S80 T71<br>T73 U75 U76 U77 U79 W72 X75 X76 X77<br>X81 Y77 Y78 | 6086    | 17.6  |

Table 2: Characteristics of the study population

|  | Characteristics of the study population (n= 345 143) |
|--|--|
| <b><u>Demographic characteristics</u></b>    |  |
| <b>Gender</b>                                |  |
| Male   | 49.5%  |
| <b>Age</b>                                   |  |
| <=12 year                                    | 14.4%  |
| 13 – 17 year                                 | 6.2%   |
| 18-25 year                                   | 9.6%   |
| 26-45 year                                   | 32.3%  |
| 46-65 year                                   | 24.7%  |
| >65 year                                     | 12.8%  |
| <b><u>Socio-economic characteristics</u></b> |  |
| <b>Highest level of education</b>            |  |
| Unknown                                      | 25.2%  |
| No education completed                       | 11.7%  |
| Primary education                            | 14.2%  |
| Secondary education                          | 36.8%  |
| Higher education                             | 12.1%  |
| <b>Health Insurance</b>                      |  |
| Unknown                                      | 23.9%  |
| Public                                       | 50.9%  |
| Private                                      | 25.3%  |
| <b>Work situation</b>                        |  |
| Work situation unknown                       | 27.9%  |
| Paid job                                     | 31.5%  |
| Attending school/ studying                   | 16.4%  |
| Housewife/ houseman                          | 11.1%  |
| Retired                                      | 9%   |
| Disability pension                           | 3%   |
| Unemployed                                   | 1.1%   |
| <b><u>Urbanicity</u></b>                     |  |
| Very strongly urban                          | 13.9%  |
| Strongly urban                               | 22.2%  |
| Moderately urban                             | 22.6%  |
| Slightly urban                               | 31.7%  |
| Non urban                                    | 9.7%   |

Table 3: the relation between having 10% more green space than average in one's living environment and the prevalence of disease clusters (n=345 143; unless stated otherwise)

| Cluster  | Percentage of green Space in 1km radius |                    | Percentage of green space in 3km radius |                    |
|--|---|--------------------|---|--------------------|
|  | OR                                      | 95% C. I.          | OR                                      | 95% C. I.          |
| <b>Cardiovascular</b>                          |   |                    |   |                    |
| High blood pressure (n = 290 535)              | 0.99                                    | 0.98 – 1.00        | 1.00                                    | 0.98 – 1.02        |
| Cardiac disease                                | 0.98                                    | 0.97 – 0.99        | 1.00                                    | 0.96 - 1.04        |
| Coronary heart disease (n = 255 346)           | <b>0.97</b>                             | <b>0.95 - 0.99</b> | 0.97                                    | 0.93 – 1.01        |
| Stroke, brain hemorrhage                       | 0.98                                    | 0.95 – 1.00        | 0.98                                    | 0.92 – 1.04        |
| <b>Musculoskeletal</b>                         |   |                    |   |                    |
| Neck- and back complaints                      | <b>0.98</b>                             | <b>0.97 - 0.99</b> | 0.99                                    | 0.97 – 1.00        |
| Severe back complaints                         | <b>0.98</b>                             | <b>0.97 - 0.99</b> | 1.00                                    | 0.98 - 1.01        |
| Severe neck and shoulder complaints            | <b>0.98</b>                             | <b>0.97 - 0.99</b> | 1.00                                    | 0.98 - 1.01        |
| Severe elbow, wrist and hand complaints        | <b>0.97</b>                             | <b>0.96 - 0.98</b> | 1.01                                    | 0.99 - 1.03        |
| Osteoarthritis (n = 255 346)                   | 0.97                                    | 0.93 - 1.01        | 0.97                                    | 0.92 – 1.03        |
| Arthritis (n = 255 346)                        | 0.99                                    | 0.97 - 1.01        | 1.00                                    | 0.96 – 1.04        |
| <b>Mental</b>                                  |   |                    |   |                    |
| Depression                                     | <b>0.96</b>                             | <b>0.95 - 0.98</b> | 0.98                                    | 0.96 – 1.00        |
| Anxiety disorder                               | <b>0.95</b>                             | <b>0.94 - 0.97</b> | <b>0.96</b>                             | <b>0.93 - 0.99</b> |
| <b>Respiratory</b>                             |   |                    |   |                    |
| Upper respiratory tract infection              | <b>0.97</b>                             | <b>0.96 - 0.98</b> | 0.99                                    | 0.97 - 1.01        |
| Bronchi(oli)tis / pneumonia                    | 0.99                                    | 0.97 – 1.00        | 1.02                                    | 0.99 – 1.04        |
| Asthma, COPD                                   | <b>0.97</b>                             | <b>0.96 - 0.98</b> | 1.01                                    | 0.99 – 1.03        |
| <b>Neurological</b>                            |   |                    |   |                    |
| Migraine/severe headache                       | <b>0.98</b>                             | <b>0.97 - 0.99</b> | 0.98                                    | 0.96 – 1.00        |
| Vertigo  | <b>0.97</b>                             | <b>0.95 - 0.99</b> | 0.98                                    | 0.94 - 1.02        |
| <b>Digestive</b>                               |   |                    |   |                    |
| Severe intestinal complaints                   | 0.98                                    | 0.96 – 1.00        | 0.99                                    | 0.95 – 1.03        |
| Infectious disease of the intestinal canal     | <b>0.97</b>                             | <b>0.95 - 0.99</b> | <b>0.95</b>                             | <b>0.91 - 0.99</b> |
| <b>Miscellaneous</b>                           |   |                    |   |                    |
| Medically Unexplained Physical Symptoms (MUPS) | <b>0.97</b>                             | <b>0.96 - 0.98</b> | <b>0.98</b>                             | <b>0.97 - 0.99</b> |
| Chronic eczema                                 | 0.99                                    | 0.97 – 1.00        | 0.99                                    | 0.95 – 1.03        |
| Acute urinary tract infection                  | <b>0.97</b>                             | <b>0.96 - 0.98</b> | 0.98                                    | 0.95 – 1.01        |
| Diabetes mellitus (n = 343 103)                | <b>0.98</b>                             | <b>0.97 - 0.99</b> | 0.98                                    | 0.97 – 1.00        |
| Cancer   | 1.00                                    | 0.98 – 1.02        | 0.99                                    | 0.95 - 1.03        |

\*Note: Odds ratio's derived from multilevel logistic regression analysis, controlling for demographic and socioeconomic characteristic and urbanicity